

## 1.5 Algebra 2.1

1. Suppose  $a, b$ , and  $c$  are positive real numbers such that  $a^2 + b^2 = c^2$  and  $\log a + \log b = \log c$ . Compute

$$\frac{(a+b+c)(a-b+c)(a+b-c)(a-b-c)}{c^2}.$$

2. Given that  $\log_{10} \sin x + \log_{10} \cos x = -1$  and that  $\log_{10} (\sin x + \cos x) = \frac{1}{2} (\log_{10} n - 1)$ , find  $n$ .

3. For some positive real number  $m$ , the acute angle  $\theta$  formed by the lines  $y = mx$  and  $my = x$  is such that  $\tan \theta = \frac{5}{12}$ . Compute  $m + 1/m$ .

4. For positive numbers  $a, b, c, d$ , show that among the four equations

$$x^2 + 2\sqrt{2a+bx} + 2\sqrt{cd} = 0,$$

$$x^2 + 2\sqrt{2b+cx} + 2\sqrt{da} = 0,$$

$$x^2 + 2\sqrt{2c+dx} + 2\sqrt{ab} = 0,$$

$$x^2 + 2\sqrt{2d+ax} + 2\sqrt{bc} = 0,$$

at least two of them each of which has two distinct real roots.

5. Graph  $y = \frac{b^x - 5}{b^x + 3}$ , assuming that the base  $b$  is greater than 1. Identify both asymptotes. Does this graph have symmetry?

## 1.9 Solving equations

1. For

$$x = \frac{4}{(\sqrt{5} + 1)(\sqrt[4]{5} + 1)(\sqrt[8]{5} + 1)(\sqrt[16]{5} + 1)},$$

compute  $(x + 1)^{48}$ .

2. Solve the equation

$$\frac{x^2 + x + 1}{x^2 + 1} + \frac{2x^2 + x + 2}{x^2 + x + 1} = \frac{19}{6}.$$

3. Let  $x_1 < x_2 < x_3$  be the three real roots of equation  $\sqrt{2014}x^3 - 4029x^2 + 2 = 0$ . Find  $x_2(x_1 + x_3)$ .

4. Find the sum of the all the real solutions to the equation

$$\sqrt{x + 2015} = x^2 - 2015.$$

5. Let  $f(x) = x^2 - 2x$ . How many distinct real numbers  $c$  satisfy  $f(f(f(f(c)))) = 3$ ?

### 4.3 Algebra 2.4

1. Let  $f(x) = x^2 + 6x + 7$ . Determine the smallest possible value of  $f(f(f(f(x))))$  over all real numbers  $x$ .
2. Compute the area of the region defined by  $x^2 + y^2 \leq |x| + |y|$ .
3. Solve the equation  $(6x + 7)^2(3x + 4)(x + 1) = 6$ .
4. Consider the parabola consisting of the points  $(x, y)$  in the real plane satisfying

$$x^2 - 2xy + y^2 - 4x + 2y + 3 = 0.$$

Find the minimum possible value of  $y$ .

5. The function  $f$ , defined by  $f(x) = \frac{ax+b}{cx+d}$ , where  $a, b, c,$  and  $d$  are nonzero real numbers, has the properties  $f(19) = 19$ ,  $f(97) = 97$ , and  $f(f(x)) = x$ , for all values of  $x$ , except  $-d/c$ . Find the range of  $f$ .